

Comparison of Renewable Natural Gas to Other Renewable Energy Technologies

March 2009

Genifuel

Overview

- **This presentation compares Catalytic Hydrothermal Gasification of Aquatic Biomass to a number of other renewable energy technologies**
- **It should be used in conjunction with the presentation titled “Renewable Natural Gas via Catalytic Hydrothermal Gasification of Aquatic Biomass”**

Compared to Biological Digestion

Biological Digestion

- **Process is well known and has been used for many years**
- **Main disadvantage is digestion time (up to 30 days) and relatively low yield**
 - Biological yield .25L methane/g biomass
 - Catalytic yield .53L methane/g biomass
- **Time in catalytic system is seconds**

Compared to High- Temperature Pyrolysis

High-Temp Gasification

- **Higher temperature and/or pressure make system design more difficult and expensive**
- **Less efficient than lower temperature catalytic system**
- **Feedstock must be dry, and must often be ground or chopped, increasing energy needs**
- **Product gas is syngas, a combination of carbon monoxide and hydrogen, which is less desirable and flexible than methane**

Compared to Algal Lipids for Diesel Fuel

Genifuel

Algal Lipids for Diesel Fuel

- **Some species of algae (and some species of diatoms) produce triacylglycerides (TAG's), in varying amounts**
- **These can be converted to a bio-diesel fuel by one of several processes:**
 - Direct hydrogenation (forming a pure alkane)
 - Transesterification (forming a fatty acid methyl ester or FAME)

Lipid Production

- **Under good growing conditions lipid production is app. 20% of cell mass**
- **To increase this to a higher level, the organism must be stressed, often by nitrogen deprivation**
 - The stress increases lipid % by weight, but essentially stops cell growth
- **Result is that high lipid content implies slow cell growth**

Lipid Production (cont.)

- **Species which produce TAGs are generally small and unicellular—often only a few microns in size**
 - These are hard to harvest
 - Small size jams filters quickly
 - Can be done by centrifugation, but this is very expensive
- **Extraction of the useful TAGs is also expensive—supercritical processes or hexane extraction requiring solvent recovery**

Lipid Production (cont.)

- **Some of the best lipid producers are diatoms, which have other difficulties**
 - Diatom shells are made of silica, which is abrasive to machinery and difficult to work with
- **Whether the chosen species are algae or diatoms, well over half of the biomass must be disposed in some way after oil extraction**
 - With algae, the general consensus is to use the remainder as cattle feed; diatoms are harder to use

Lipid Production Economics

- **The combination of slow growth, expensive harvesting and oil extraction, and relatively low yield (less than half of the mass) makes this an expensive fuel source**
- **Unlikely to be profitable without subsidies in the foreseeable future (10 years?)**

Compared to Oil Crops

Fuel from Oil Crops

- **Many food-type crops produce vegetable oils which can be converted to bio-diesel fuel**
 - Soybeans, canola, peanuts, palm, sunflower, etc.
- **The problem with using food crops is that it creates competition between food and fuel for premium land and water (similar to corn or sugar ethanol)**
- **This competition is not desirable or economic in the long run**

Compared to Corn or Sugar Ethanol

Genifuel

Corn Ethanol

- **In the US, by far the largest volume of current biofuel production is ethanol from corn**
- **Corn is grown by conventional means, and the grain is processed to remove the starch, which is fermented to alcohol**
- **The technology is well known, proven, and efficient.**

Sugar Ethanol

- **Ethanol from sugar is the dominant biofuel in Brazil, and one of the major biofuel products in the world**
- **However, it suffers from many of the same problems as corn ethanol**
- **Sugar ethanol is unlikely to be economically viable in the long run**

Corn Ethanol (cont.)

- **Using corn for fuel has highlighted a number of problems**
 - Corn production is energy intensive (for nitrogen fertilizer and other agricultural requirements)
 - Corn requires abundant water and good land
 - Corn is a basic food grain, and using corn for fuel creates competition between food and fuel
 - The net result is that corn ethanol is only slightly energy positive, and is too expensive in the long run

Compared to Cellulosic Ethanol

Genifuel

Cellulosic Ethanol

- **Cellulosic ethanol production is generally assumed to come from terrestrial biomass**
- **The process involves the thermal, chemical, or biological breakdown of plant material to extract sugars**
- **The sugars are then fermented biologically into ethanol**
- **The alcohol can be used as fuel or blended with hydrocarbon fuels such as gasoline**
- **Not clear how to use remaining material**

Cellulosic Ethanol

- **Cellulosic ethanol is an alternative to fuel production from food crops**
- **Current research focuses on fast-growing species which can use marginal (non-crop) land**
 - Examples include miscanthus, switchgrass, etc.

Cellulosic Ethanol (Problems)

- **Current production is non-economic, primarily because extracting sugars is still technically difficult and expensive**
- **Very high yields (30 tons/acre/yr) are often quoted for crops such as miscanthus or switchgrass, but such yields can only be achieved with ample water and fertilizer.**
- **On marginal land, with no irrigation or fertilizer, the yields are likely to be in the range of 5-7 tons/acre or less**

Compared to Photovoltaics (Solar Cells)

Photovoltaic vs. Biological Systems

- **Photovoltaic energy production is very similar to growth of biomass in the sense that both depend on capturing photons from sunlight**
- **PV systems produce electricity directly for immediate use, while biological systems store energy in chemical form for later use**
- **Both capture similar energy per area of land**

Photovoltaic vs Biological Systems (cont)

- **Both accomplish a quantum conversion of photons to usable energy**
 - PV systems use semiconductor materials to achieve charge separation (electrons and holes)
 - Biological systems use photosynthesis to achieve charge separation as electrons and hydrogen ions (protons)
- **Both systems use a similar portion of the solar spectrum and (with current technology) achieve similar quantum efficiencies**

Major Differences

- **PV and Biological systems have two major differences:**
 - Energy storage
 - PV energy output is electricity which must be used immediately
 - Biological energy can be harvested, stored, and converted to usable energy day or night
 - After conversion to renewable natural gas, the energy (in the form of fuel) can be stored indefinitely without loss
 - PV systems are vastly more capital intensive—by at least an order of magnitude

PV Economics

- **The high cost (capital intensity) of PV systems make them uneconomic today**
- **Typical of semiconductor technology, PV costs are declining**
- **Most forecasts expect economic viability around 2015**

Major Differences

- **PV systems produce peak output from late morning to mid-afternoon**
- **Renewable natural gas from aquatic biomass can be used on demand, including by conversion to electricity on demand via natural gas turbines**
 - Can be used as “peaking power” when needed
 - Can be used as “base load” power, which is not possible with PV power

Compared to Wind Turbines

Genifuel

Wind Turbines

- **Wind turbines are usable only in areas of moderate but consistent wind**
- **Fortunately, the US has a number of such areas**
- **The problem is that good wind areas are usually in remote locations with no connection to the electrical grid**
- **High concentrations of wind turbines can also be noisy and unsightly, further driving them to remote locations**

Wind Turbines (cont)

- **Wind turbines are capital intensive compared to biological growth systems—wind is similar to PV systems in capital intensity**
- **Wind systems achieve relatively low capital utilization—often in the range of 20% to 30%**
- **Much confusion arises because nameplate capacities are not the same as effective capacities—the only valid measure of wind capacity is total energy produced per year, not nameplate capacity of the turbines**

Compared to Concentrated Solar

Genifuel

Concentrated Solar

- **Concentrated solar uses mirrors to focus sunlight on a thermal collector**
- **Heat is generally converted to steam to produce electricity with a conventional steam turbine-generator**
- **Process heat may also be made**
- **Economics are similar to wind**

Conclusion

Conclusion

- **Wind, PV, Concentrated Solar, and Biomass Gasification all achieve a similar power density—that is, the amount of power which can be produced per unit of land**
- **This figure is in the range of 750-850 MWh/acre/year**
- **Biomass gasification has the lowest capital cost, and is the only one of these technologies which can provide continuous energy production 24/7**

Conclusion (cont.)

- **At the current stage of development, the production of renewable natural gas via catalytic hydrothermal gasification of aquatic biomass is very likely the most economically viable renewable energy process**
- **Genifuel produces renewable natural gas via this process**